

$N-\bar{N}$ Working Group Summary

Transformative Hadron Beamlines Workshop

BNL July 21-23, 2014

Mary Bishai (BNL), Gustaaf Brooijmans (Columbia U.)

July 23, 2014

- 1 $N - \bar{N}$ Goals
- 2 Physics Motivation
- 3 $n - \bar{n}(\prime)$ Oscillation Experimental Concepts
- 4 R&D for Spallation Sources
- 5 Proton Radiography
- 6 Summary of $N - \bar{N}$ Needs

$N - \bar{N}$ at THB Workshop

$N - \bar{N}$ Working Group Summary

Mary Bishai
(BNL),
Gustaaf Brooijmans
(Columbia U.)

$N - \bar{N}$ Goals

Physics
Motivation

$n - \bar{n}(\prime)$
Oscillation
Experimental
Concepts

R&D for
Spallation
Sources

Proton
Radiography

Summary of
 $N - \bar{N}$ Needs

How We Got Here

- ❖ Collaboration is being formed to pursue new search for neutron-anti-neutron oscillations using intense ESS cold neutron flux
 - Expect to improve sensitivity by $\mathcal{O}(500)$
 - www.nnbar-at-ess.org
- ❖ Question arose: what can be done at BNL?
 - Not a competitive oscillation experiment
 - At least not horizontal
 - (Semi-)vertical??
 - Discuss this over coffee or beer
 - Neutron test beams?
 - Other things “NNbar enthusiasts” are interested in?



$n - \bar{n}$ Oscillations Beyond the Standard Model

$N - \bar{N}$ Working
Group
Summary

Mary Bishai
(BNL),
Gustaaf
Brooijmans
(Columbia U.)

$N - \bar{N}$ Goals

Physics
Motivation

$n - \bar{n}(\prime)$
Oscillation
Experimental
Concepts

R&D for
Spallation
Sources

Proton
Radiography


Summary of
 $N - \bar{N}$ Needs

- Many reasons to expect B-number is **not a good symmetry** of nature
- **Sphalerons in SM, GUTs, origin of matter, etc...**
- If B is violated, important to determine the selection rules:
- **$\Delta B = 1$ (proton decay)**
- **$\Delta B = 2$ (neutron-antineutron oscillations)**
- For many extensions of the standard model (Pati-Salam models with supersymmetry, theories with extra dimensions or branes, model-independent treatments)

$N - \bar{N}$ can occur at experimental limits even when proton decay not observed!

Is There Mirror Matter?


Two coexisting worlds look theoretically very attractive



- Two identical gauge factors, $G \times G'$, with identical field contents and Lagrangians: $\mathcal{L}_{\text{tot}} = \mathcal{L} + \mathcal{L}' + \mathcal{L}_{\text{mix}} \quad - \quad SU(5) \times SU(5)', \text{ etc.}$
- Can naturally emerge in string theory: O & M matter fields localized on two parallel branes with gravity propagating in bulk: e.g. $E_8 \times E'_8$
- Exact parity $G \leftrightarrow G'$: Mirror matter is dark (for us), but its particle physics we know exactly (on our skin) – **no new parameters!**

Next Generation $n - \bar{n}$ Oscillation Search Expts (R. Pattie)

Horizontal Configuration for Next Generation NNbar search experiment



1. Increased flight path
2. Colder neutron source
3. Higher m supermirror neutron optics
4. Modern Calorimeter/Vertex Tracker

Challenges of Nex-Gen $n - \bar{n}$ Oscillation Expts

$N - \bar{N}$ Working
Group
Summary

Mary Bishai
(BNL),
Gustaaf
Brooijmans
(Columbia U.)

$N - \bar{N}$ Goals

Physics
Motivation

$n - \bar{n}(\prime)$
Oscillation
Experimental
Concepts

R&D for
Spallation
Sources


Proton
Radiography

Summary of
 $N - \bar{N}$ Needs

- Previous generation of $n - \bar{n}$ oscillation expts at reactors. Nex-Gen is at accelerators (spallation sources) \Rightarrow more backgrounds from fast neutrons and protons.
- Need better detectors to constrain annihilation vertex \Rightarrow more tests of detector technology in neutron beams to understand fast background rejection.
- Pulsed beams can also help reduce backgrounds.
- ATLAS TRT tests at LANL LANSCE used 5-10n/seconds. Factor of 10x intensity less could still be useful to benchmark detector simulations of the fast neutron background.
- A big effort, $n - \bar{n}$ oscillations is not within the scope of what can be done at BNL.
BNL can participate in detector and moderator R&D.

Mirror Matter $n - n'$ Oscillation Searches (Y. Kamyshev)

Neutron oscillating into mirror neutron is interacting with the trap wall



in case of successful guessing for $\vec{B} = \vec{B}'$
the resonance enhancement is expected: the
oscillation frequency will be reduced to (1/few s)
and oscillation amplitude increased by
few orders of magnitude, ultimately to

$$P_{nn'} = \left(\frac{t}{\tau_{nn'}} \right)^2$$

t — observation time
 $\tau_{nn'}$ — oscillation time
 > 1 sec

A signal for mirror matter?

$N - \bar{N}$ Working
Group
Summary

Mary Bishai
(BNL),
Gustaaf
Brooijmans
(Columbia U.)

$N - \bar{N}$ Goals

Physics
Motivation

$n - \bar{n}(\prime)$
Oscillation
Experimental
Concepts

R&D for
Spallation
Sources

Proton
Radiography

Summary of
 $N - \bar{N}$ Needs

**Magnetic anomaly in UCN
trapping: signal for neutron
oscillation to parallel world?**

Z. Berezghiani and F. Nesti
Eur. Phys. J. C72 (2012) 1974;
also <http://arxiv.org/abs/1203.1035>

$$A_B^{\text{det}}(t) = \frac{N_{-B}(t) - N_B(t)}{N_{-B}(t) + N_B(t)}$$

Measured asymmetry \rightarrow
 $\sim (7 \pm 1.4) \times 10^{-4}$ ($\sim 5\sigma$)





Fig. 1. Upper Panel: from up to down, the monitor and detector counts in $\{B\}$ series, M and $N = N_1 + N_2$ normalized respectively to 470000 and 140000; and the ratios $N/M (\times 47/14)$ and N_1/N_2 . Lower Panel: results for A_B^{det} binned by two $\{B\}$ cycles (16 measurements), with the constant and periodic fits.

New concept: Neutron Regeneration (Y. Kamyshev)

Possible Neutron Regeneration Search

- It is an **appearance** search of $n \rightarrow n' \rightarrow n$
- Alternative to **disappearance** $n \rightarrow n'$ observed with UCN
- It excludes collisions with walls that might be the reason for some unknown effect resulting to the measured asymmetry with UCN.



Mirror Matter Search at BNL?

$N - \bar{N}$ Working
Group
Summary

Mary Bishai
(BNL),
Gustaaf
Brooijmans
(Columbia U.)

$N - \bar{N}$ Goals

Physics
Motivation

$n - \bar{n}(\prime)$
Oscillation
Experimental
Concepts

R&D for
Spallation
Sources

Proton
Radiography





Summary of
 $N - \bar{N}$ Needs

- Small experiment that could be carried out at BNL.
- A BNL $n \rightarrow n' \rightarrow n$ regeneration experiment is complementary to $n \rightarrow n'$ disappearance experiment done earlier with a potential 5.2σ signal.
- Could be done now with 100kW for 1 day or 20 kW for 10 days for example.
- Requires cold neutrons
- Pulsed beam structure can be used to reduce backgrounds.

ESS Moderator Designs (G. Muhrer)



TDR configuration: two tall moderators



Thermal wings provide a bi-spectral source.

VOLUME moderator:
implemented at J-
PARC

Flat Vs. Tall Moderators (G. Muhrer)

$N - \bar{N}$ Working
Group
Summary

Mary Bishai
(BNL),
Gustaaf
Broijmans
(Columbia U.)

$N - \bar{N}$ Goals




Physics
Motivation

$n - \bar{n}(\prime)$
Oscillation
Experimental
Concepts

R&D for
Spallation
Sources

Proton
Radiography

Summary of
 $N - \bar{N}$ Needs




□ thermal neutrons arriving from the surroundings are transformed into cold ones within about 1 cm of the walls of the moderator vessel

ESS Flat Moderator Concept (G. Muhrer)



Flat moderator reference configuration



Upper moderator view from above
Proton beam comes from the left

MR plug view in proton beam direction
Proton beam comes from behind

$N - \bar{N}$ Working
Group
Summary

Mary Bishai
(BNL),
Gustaaf
Brooijmans
(Columbia U.)

$N - \bar{N}$ Goals

Physics
Motivation

$n - \bar{n}(\prime)$
Oscillation
Experimental
Concepts


R&D for
Spallation
Sources

Proton
Radiography

Summary of
 $N - \bar{N}$ Needs

Bottom Moderator Designs (G. Muhrer)

High intensity D₂ moderator



High intensity D₂ source can give neutron intensity (brightness × emission surface area) **3-4 × TDR**

Moderator tests at BNL THB?

$N-\bar{N}$ Working
Group
Summary

Mary Bishai
(BNL),
Gustaaf
Brooijmans
(Columbia U.)

$N - \bar{N}$ Goals

Physics
Motivation

$n - \bar{n}(\prime)$
Oscillation
Experimental
Concepts

R&D for
Spallation
Sources

Proton
Radiography

Summary of
 $N - \bar{N}$ Needs

- ESS is going forward
- Possible BNL contribution to experiments using cold neutrons from ESS:
test new moderator materials (designs?) in a test facility.
- Need 100 kW (?)test facility where we can test an engineered model in an environment where the degradation of the moderator can be monitored in a low intensity environment.
- Most likely its the material of the bottom moderator (could serve up to 11 beamlines) that can be subject to R&D - like D_2 .

Proton Radiography (A. Saunders)

$N - \bar{N}$ Working
Group
Summary

Mary Bishai
(BNL),
Gustaaf
Brooijmans
(Columbia U.)

$N - \bar{N}$ Goals

Physics
Motivation

$n - \bar{n}(\prime)$
Oscillation
Experimental
Concepts



R&D for
Spallation
Sources

Proton
Radiography

Summary of
 $N - \bar{N}$ Needs

Different probes- 1×10^9 incident particles.

French test object (FTO)




bb's
 $\lambda = 0. \text{ g/cm}^2$

X-rays
 $\lambda = 25. \text{ g/cm}^2$

Protons
 $\lambda \approx 185. \text{ g/cm}^2$

Neutrinos
 $\lambda = \infty \text{ g/cm}^2$

Simulated radiograph Density reconstruction




Proton Radiography (A. Saunders)

Resolution of Proton Radiography


1. **Object scattering** - introduced as the protons are scattered while traversing the object.
2. **Chromatic aberrations**- introduced as the protons pass through the magnetic lens imaging system.
3. **Detector blur**- introduced as the proton interacts with the proton-to-light converter and as the light is gated and collected with a camera system.

Object Scattering



$$\sigma_o = \frac{1}{\sqrt{3}} \theta \frac{l}{2} = \frac{14.1}{\sqrt{6}} \frac{1}{P\beta} \sqrt{\frac{l^2}{x_o^2} \propto \frac{l^2}{P^2}}$$


Chromatic Aberrations



$$\sigma_c = l_c \theta \frac{\delta P}{P} = c \sqrt{P} \frac{\delta P}{P^2} \frac{14.1}{\beta} \sqrt{\frac{l}{x_o} \propto \sqrt{\frac{l}{P^3}}}$$

Assume detector development can keep up

Detector Blur



$$\sigma_s = \theta_s \frac{l_s \sqrt{l}}{P}$$

Scintillator

Camera

Resolution is independent of proton energy

Previous Radiography Facility at AGS (A. Saunders)

N- \bar{N} Working Group Summary

Mary Bishai (BNL), Gustaaf Brooijmans (Columbia U.)

N — \bar{N} Goals

Physics Motivation

$n - \bar{n}(\prime)$
Oscillation
Experimental Concepts

R&D for Spallation Sources

Proton Radiography

Summary of *N* — \bar{N} Needs

Lens system and camera station for 24 GeV radiography at the AGS (Experiment 933)


24 GeV
 ~ 30 ns pulses
 $< 1 \times 10^{11}$ protons/pulse

pRad lens in U-Line
at AGS



P Radiography Industrial Applications (A. Saunders)

Industrial Applications could take advantage of NNSA-funded pRad facility



Vacuum Pump Turbine
(Molecular Drag pump)

Engine Head

Possible Radiography Facilities at BNL (A. Saunders)

$N - \bar{N}$ Working
Group
Summary

Mary Bishai
(BNL),
Gustaaf
Brooijmans
(Columbia U.)

$N - \bar{N}$ Goals

Physics
Motivation

$n - \bar{n}(\prime)$
Oscillation
Experimental
Concepts

R&D for
Spallation
Sources

Proton
Radiography

Summary of
 $N - \bar{N}$ Needs



Summary of Beam Requirements

$N-\bar{N}$ Working
Group
Summary

Mary Bishai
(BNL),
Gustaaf
Brooijmans
(Columbia U.)

$N - \bar{N}$ Goals

Physics
Motivation

$n - \bar{n}(\prime)$
Oscillation
Experimental
Concepts

R&D for
Spallation
Sources

Proton
Radiography

Summary of
 $N - \bar{N}$ Needs

Species	Beam Energy	Intensity	Rep Rate	Custom	Comment
p	0.5-2	high	$O(10)$		n moderator testing close to spallation target
n	cold (few meV)			$\sim 10^8$ n/s	neutron regeneration
n	fast (10-800 MeV)			10^6 n/s?	WNR alternative
p	24 GeV	$1e11$			p radiography only few pulses needed